

THE SIMPLE CLASSICAL GROUPS OF DIMENSION LESS THAN 6 WHICH ARE $(2, 3)$ -GENERATED

M.A. PELLEGRINI AND M.C. TAMBURINI BELLANI

ABSTRACT. In this paper we determine the classical simple groups of dimension $n = 3, 5$ which are $(2, 3)$ -generated (the cases $n = 2, 4$ are known). If $n = 3$, they are $\mathrm{PSL}_3(q)$, $q \neq 4$, and $\mathrm{PSU}_3(q^2)$, $q^2 \neq 9, 25$. If $n = 5$ they are $\mathrm{PSL}_5(q)$, for all q , and $\mathrm{PSU}_5(q^2)$, $q^2 \geq 9$. Also, the soluble group $\mathrm{PSU}_3(4)$ is not $(2, 3)$ -generated. We give explicit $(2, 3)$ -generators of the linear preimages, in the special linear groups, of the $(2, 3)$ -generated simple groups.

1. INTRODUCTION

A $(2, 3)$ -generated group is a group that can be generated by two elements of respective orders 2 and 3. Alternatively, by a result of Fricke and Klein [5], it can be defined as an epimorphic image, of order > 3 , of the group $\mathrm{PSL}_2(\mathbb{Z})$.

Many finite simple and quasi-simple groups have been investigated with respect to this property. Apart from the infinite families $\mathrm{PSP}_4(2^m)$ and $\mathrm{PSp}_4(3^m)$, the other finite classical simple groups are $(2, 3)$ -generated, up to a finite number of exceptions, by the probabilistic results of Liebeck and Shalev [8]. These exceptions, for groups of Lie type, occur in small dimensions over small fields. For a recent survey see [16]. Our notation for classical groups is taken in accordance with [2]: consequently we denote by $\mathrm{SU}_n(q^2)$ the group appearing also as $\mathrm{SU}_n(q)$ in the literature, e.g. in [1].

The groups $\mathrm{PSL}_2(q)$ were dealt with by Macbeath in [9]: the only exception is $\mathrm{PSL}_2(9)$. The groups $\mathrm{SL}_4(q)$ and $\mathrm{SU}_4(q^2)$ were studied in [11]. In dimension 4, apart from the above infinite series of symplectic groups, the only simple groups which are not $(2, 3)$ -generated are $\mathrm{PSL}_4(2)$, $\mathrm{PSU}_4(4)$, $\mathrm{PSU}_4(9)$. The groups $\mathrm{SL}_n(q)$, for $n = 5, 6, 7$ were studied in [4], [14], [13], [12]: all these groups are $(2, 3)$ -generated. The aim of this paper is to fill the gaps for classical groups of dimension $n \leq 5$, treating in a uniform way the special linear and the unitary groups. This also gives a partial answer to Problem n. 18.98 of [10].

The soluble group $\mathrm{PSU}_3(4)$ and the simple groups $\mathrm{PSL}_3(4)$ (Garbe [6], Cohen [3]) and $\mathrm{PSU}_3(9)$ (Wagner [17]) are not $(2, 3)$ -generated. The same applies to $\mathrm{PSU}_3(25)$ by Theorem 3.6 of this paper. In Theorems 3.5 and 3.7 we show that:

- (i) $\mathrm{SL}_3(q)$ and $\mathrm{PSL}_3(q)$ are $(2, 3)$ -generated for all $q \neq 4$;
- (ii) $\mathrm{SU}_3(q^2)$ and $\mathrm{PSU}_3(q^2)$ are $(2, 3)$ -generated for all $q^2 \neq 4, 9, 25$.

The group $\mathrm{PSU}_5(4)$ is not $(2, 3)$ -generated (Vsemirnov [15]). On the other hand, in Theorems 4.6 and 4.7 we show that:

- (i) $\mathrm{SL}_5(q)$ and $\mathrm{PSL}_5(q)$ are $(2, 3)$ -generated for all q ;

2010 *Mathematics Subject Classification.* 20G40, 20F05.

Key words and phrases. Generation; Classical groups.

(ii) $\mathrm{SU}_5(q^2)$ and $\mathrm{PSU}_5(q^2)$ are $(2, 3)$ -generated for all $q^2 \geq 9$.

The proofs of our results are based on the classification of maximal subgroups of the finite classical groups ([1]) which, in the Aschbacher notation, fall into classes $\mathcal{C}_1, \dots, \mathcal{C}_8$ and $\mathcal{S}$.

2. NOTATION AND PRELIMINARY RESULTS

Let $\mathbb{F}$ be an algebraically closed field of characteristic $p \geq 0$. When $p > 0$, we set $q = p^m$, $m \geq 1$. We denote by σ the automorphism of $\mathrm{SL}_n(q^2)$ defined as:

$$(1) \quad (\alpha_{i,j}) \mapsto (\alpha_{i,j}^q).$$

For a fixed n , we call x, y two elements of $\mathrm{SL}_n(\mathbb{F})$ whose respective orders are 2 and 3. We denote by z the product xy , and by H the subgroup of $\mathrm{SL}_n(\mathbb{F})$ generated by x, y , i.e., $H = \langle x, y \rangle$.

We need H to be absolutely irreducible. This requirement restricts the possible canonical forms of x, y and their shapes, up to conjugation.

We fix certain roots of unity in $\mathbb{F}^*$, according to the notation in Table A.

Table A

Root	Order	Root	Order	Root	Order
ω	$\frac{3}{(3,p)}$	i	$\frac{4}{(4,p^2)}$	η	$\frac{5}{(5,p)}$

For later use, we give some lower bounds of the Euler-Phi function $\varphi : \mathbb{N} \rightarrow \mathbb{N}$.

Lemma 2.1. *Let n be a natural number. Then $\varphi(n) > n^{2/3}$ if, and only if,*

$$n \notin \{1, 2, 3, 4, 6, 8, 10, 12, 18, 24, 30, 42\}.$$

Furthermore, $\varphi(n) = n^{2/3}$ if, and only if, $n = 1, 8$.

Proof. Recall that if m, n are two coprime integers, then $\varphi(mn) = \varphi(m)\varphi(n)$. Let us first consider the case $n = s^a$, with s a prime. Then

$$\varphi(s^a) = s^{a-1}(s-1) > (s^a)^{2/3} \Leftrightarrow s^{a-3}(s-1)^3 > 1.$$

The last inequality holds if, and only if, one of the following cases occurs:

(i) $a = 1$, $s \neq 2, 3$; (ii) $a = 2, 3$, $s \neq 2$; (iii) $a \geq 4$.

In particular this implies that, if $(n, 6) = 1$, then $\varphi(n) > n^{2/3}$. So, we are left to deal with the cases when $(n, 6) \neq 1$. Suppose first $n = 2^a 3^b$ with $a, b \geq 1$. Then

$$\varphi(2^a 3^b) > (2^a 3^b)^{2/3} \Leftrightarrow (2^a 3^b) > 27.$$

The last inequality holds if, and only if, one of the following occurs:

(i) $b = 1$ and $a \geq 4$; (ii) $b = 2$ and $a \geq 2$; (iii) $b \geq 3$ and $a \geq 1$.

Finally, let $n = uv$, where $(u, v) = 1$ and $u = 2^a 3^b$, with $a + b \geq 1$. By the above, if $u \notin E := \{2, 3, 4, 6, 8, 12, 18, 24\}$, then $\varphi(n) > n^{2/3}$. On the other hand, if $u \in E$, then $\varphi(u) \geq \frac{3}{5}u^{2/3}$. Thus, it suffices to prove that $\varphi(v) > \frac{5}{3}v^{2/3}$. Note that, for all primes $s \geq 11$, $\varphi(s) > \frac{5}{3}s^{2/3}$ and so $\varphi(s^a) > s^{2(a-1)/3}\varphi(s) > \frac{5}{3}(s^a)^{2/3}$. For the exceptional cases $s = 5, 7$, we have that $\varphi(s^a) > \frac{5}{3}(s^a)^{2/3}$ if, and only if, $a \geq 2$. Also, observe that $\varphi(35) > \frac{5}{3} \cdot (35)^{2/3}$. Hence, we are left to consider only the integers $n = uv$, where $u \in E$ and $v \in \{1, 5, 7\}$. Direct computations lead to the exceptions listed in the statement. $\square$

The following corollary will be used for $n = q = p^m$.

Corollary 2.2. *For all $n \geq 14$, we have $\varphi(n^2 - 1) > \max\{3n + 21, 4n - 1\}$.*

Proof. If $n \geq 64$, we may apply Lemma 2.1. In fact, for these values of n , it is easy to see that $(n^2 - 1)^2 > (3n + 21)^3$ and $(n^2 - 1)^2 > (4n - 1)^3$. So $\varphi(n^2 - 1) > \max\{3n + 21, 4n - 1\}$. For the values $14 \leq n \leq 63$ we deduce this inequality from direct computations. $\square$

In order to exclude that $H = \langle x, y \rangle$ is contained in a maximal subgroup of class $\mathcal{C}_6$ or class $\mathcal{C}_3$ we will use respectively Lemma 2.3 and Lemma 2.4.

Lemma 2.3. *Let $x, y \in \mathrm{SL}_r(q)$, with r an odd prime. Assume that $x^2 = y^3 = 1$. If $(xy)^6$ is not scalar, the projective image $\overline{H}$ of H is not contained in any subgroup $\overline{M}$ of $\mathrm{PSL}_r(q)$ having the following structure: $\overline{M} = N.S$, where N is a normal elementary abelian subgroup of order r^2 , and $S \leq \mathrm{SL}_2(r) = \mathrm{Sp}_2(r)$ acts by conjugation on N , according to the natural action of $\mathrm{SL}_2(r)$ on $\mathbb{F}_r^2$.*

Proof. Assume, by contradiction, that $\overline{H} \leq \overline{M}$. Since r is odd, the only involution of $\mathrm{SL}_2(r)$ is the central one, namely $-I$. Thus Nx is central in $\overline{M}/N \simeq S$. It follows $N(xy)^3 = Nx^3y^3 = Nx$, i.e., $(xy)^3x^{-1} = n_1 \in N$. Moreover, recalling from the statement that the action of Nx on $N \simeq \mathbb{F}_r^2$ is defined by $n \mapsto x^{-1}nx$, we get $x^{-1}nx = n^{-1}$, for all $n \in N$. It follows $(xy)^3 = n_1x$, with $n_1 \in N$, whence $(xy)^6 = n_1xn_1x = n_1n_1^{-1} = 1$, in contrast with the assumption that $(xy)^6$ is not scalar. $\square$

Lemma 2.4. *For $q = p^m$, $m > 1$, let $a \in \mathbb{F}_q^*$ have order $q - 1$. For all primes s*

$$\mathbb{F}_p[a^s] = \mathbb{F}_q$$

except, possibly, when $s = \sum_{j=0}^{t-1} q_0^j$, with $q = q_0^t$ for some $t \geq 2$.

In particular the statement is true for $s = 3$, $q \neq 4$ and for $s = 5$, $q \neq 16$.

Proof. If $(s, q - 1) = 1$ our claim is obvious. So we may assume that $q - 1 = sk$. Suppose $\mathbb{F}_p[a^s] = \mathbb{F}_{q_0}$ for some q_0 such that $q = q_0^t$ with $t \geq 2$. Since k is the order of a^s , we have $q_0 - 1 \geq k$, whence:

$$sk = q - 1 = (q_0 - 1) \left(\sum_{j=0}^{t-1} q_0^j \right) \geq k \left(\sum_{j=0}^{t-1} q_0^j \right).$$

This gives $s \geq \sum_{j=0}^{t-1} q_0^j$. If s divides $q_0 - 1$ we get the contradiction

$$\sum_{j=0}^{t-1} q_0^j \leq s \leq q_0 - 1.$$

Thus s , being a prime, must divide $\sum_{j=0}^{t-1} q_0^j$, whence $s = \sum_{j=0}^{t-1} q_0^j$. $\square$

3. DIMENSION 3

Let X, y be elements of $\mathrm{SL}_3(\mathbb{F})$ such that their projective images have orders 2, 3 and generate an absolutely irreducible subgroup H . Then, $X = \omega^j x$, for some $j = 0, 1, 2$, with $x^2 = I$. We may replace X by x , leaving unchanged the projective image of H . The involution x must be conjugate to blockdiag $\left(-1, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right)$. Hence x has a 2-dimensional eigenspace W , relative to the eigenvalue -1 . Clearly $W \cap yW$ has dimension 1. So we may assume $W = \langle e_1, e_2 \rangle$ with $e_2 = ye_1$, and $ye_2 \notin W$. Setting $ye_2 = e_3$, it follows $ye_3 = \lambda e_1$. The condition $\det(y) = 1$ gives

$\lambda = 1$. In particular $y^3 = I$. Set $xe_3 = ae_1 + be_2 + e_3$ with $a, b \in \mathbb{F}$. For $p = 2$, we need $(a, b) \neq (0, 0)$ in order that x is an involution. For p odd and $(a, b) = (0, 0)$ the group H is isomorphic to $\text{Alt}(4)$. Hence, for our purposes, we may assume:

$$(2) \quad x = \begin{pmatrix} -1 & 0 & a \\ 0 & -1 & b \\ 0 & 0 & 1 \end{pmatrix}, \quad (a, b) \neq (0, 0), \quad y = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

Lemma 3.1. *Let x, y be defined as in (2). Then H is absolutely irreducible if, and only if, the following conditions hold. For all $j \in \{0, 1, 2\}$:*

- (i) $p \neq 2$ and $(a, b) \neq (2\omega^j, 2\omega^{2j})$;
- (ii) $b \neq -a\omega^j - 2\omega^{2j}$.

Proof. H fixes some 1-dimensional space precisely when $p \neq 2$ and $(a, b) = (2\omega^j, 2\omega^{2j})$, for some $j = 0, 1, 2$. In this case, setting $u = (a, b, 2)^T$ we have $u \neq 0$ and $xu = u, yu \in \langle u \rangle$.

H fixes a 2-dimensional space if, and only if, H^T fixes a 1-dimensional space. This happens only if $b = -a\omega^j - 2\omega^{2j}$, for some $j = 0, 1, 2$. In this case, taking $w = (1, \omega^{2j}, \omega^j)^T$, we have $x^T w = -w$ and $y^T w = \omega^{2j} w$. $\square$

Lemma 3.2. *Assume that H is absolutely irreducible and that $ab \neq 1$. Then H is not conjugate to a subgroup of the monomial group.*

Proof. Assume that H acts monomially on a basis $B = \{v_1, v_2, v_3\}$. We may suppose $xv_1 \in \langle v_1 \rangle$ and, furthermore, $xv_1 = -v_1$. The latter claim is clear if x fixes all the subspaces $\langle v_j \rangle$; it follows from $x^2 = I$ and the determinant condition, if x interchanges $\langle v_2 \rangle$ with $\langle v_3 \rangle$. Thus $v_1 = (c, d, 0)^T$ for some $(c, d) \neq (0, 0)$. By the irreducibility of H , we may assume $v_2 = yv_1 = (0, c, d)^T, v_3 = yv_2 = (d, 0, c)^T$. The condition $(a, b) \neq (0, 0)$ implies that B cannot be the canonical basis (up to scalars). Thus $cd \neq 0$: in particular x cannot fix $\langle v_2 \rangle$ and $\langle v_3 \rangle$. From $xv_2 = (ad, -c + bd, d)^T$ we see that $xv_2 \in \langle v_3 \rangle$ if, and only if, $ab = 1$. $\square$

Lemma 3.3. *Assume that H is absolutely irreducible and that $ab \neq 1$. Set $z = xy$.*

- (i) z^j is not scalar for $j \leq 4$ and for $j = 6$.
- (ii) If z^5 is scalar then both a and b are roots of $t^6 - 4t^3 - 1$.

More precisely:

- if $p = 2$, then $(a, b) \in \{(\omega^j, 1), (1, \omega^j), (\omega^j, \omega^j)\}$, for some $j = 1, 2$;
- if $p \neq 2$, then $b = \frac{1 \pm \sqrt{5}}{2} \omega^j, a = b\omega^j$ for some $j = 0, 1, 2$.
- (iii) If z^7 is scalar then:
 - either $p = 2$ and $(a, b) \in \{(\omega^j, 0), (0, \omega^j)\}$, for some $j \in \{0, 1, 2\}$;
 - or $a \neq 0, b \neq 0$ and both a and b are roots of

$$R = t^{15} - 16t^{12} + 59t^9 - 67t^6 - 37t^3 + 8.$$

R splits over a field containing an element of order $\frac{21}{(p, 21)}$ and factorizes over $\mathbb{Z}$ as

$$(t^2 + t + 2)(t^3 - 2t^2 - t + 1)(t^4 - t^3 - t^2 - 2t + 4)(t^6 + 2t^5 + 5t^4 + 3t^2 + t + 1).$$

Proof. We consider the action of z on the canonical basis $\{e_1, e_2, e_3\}$.

(i) For $j \leq 4$ our claim follows easily from the assumptions, since:

$$\begin{aligned} ze_3 &= -e_1, & z^2e_3 &= e_2, \\ z^3e_3 &= ae_1 + be_2 + e_3, & z^4e_3 &= (ab - 1)e_1 + (-a + b^2)e_2 + be_3. \end{aligned}$$

Assume z^6 to be scalar and note that $z^6e_3 = f_1e_1 + f_2e_2 + f_3e_3$, where

$$f_1 = -2a^2b + ab^3 + 2a - b^2, \quad f_2 = a^2 - 3ab^2 + b^4 + 2b.$$

Clearly $b \neq 0$ as $b = 0$ would give also $a = 0$. From $f_1 = (2a - b^2)(1 - ab)$, we get $2a = b^2$ as we are excluding $ab = 1$. So $p \neq 2$ and $a = \frac{b^2}{2}$. Substitution in f_2 gives $b^4 = 8b$. It follows that $(a, b) = (2\omega^j, 2\omega^{2j})$, against the assumption that H is absolutely irreducible.

(ii) Assume z^5 to be scalar and note that $z^5e_3 = f_1e_1 + f_2e_2 + f_3e_3$ where

$$f_1 = -a^2 + ab^2 - b, \quad f_2 = -2ab + b^3 + 1.$$

Notice that $b \neq 0$, since $b = 0$ would give $a = 0$. Similarly $a \neq 0$. Considering f_1 and f_2 as polynomials in a , their resultant $\text{Res}(f_1, f_2)$ is $b^2(b^6 - 4b^3 - 1)$. By [7, Theorem 5.7, p. 325], where the definition and the properties of resultants are given, the non-zero values of b satisfying $f_1 = f_2 = 0$ must be roots of

$$R = t^6 - 4t^3 - 1 = (t^2 - t - 1)(t^2 - \omega t - \omega^2)(t^2 - \omega^2 t - \omega).$$

Calling t_1, t_2 the roots of $t^2 - t - 1$, the roots of P are $t_k\omega^j$, $k = 1, 2$, $j = 0, 1, 2$. Since b is a root of P , we may set $b = t_k\omega^j$, whence $b^3 + 1 = 2(t_k + 1)$. Now we equate to zero the coefficient of e_2 . If $p = 2$, we may assume $t_1 = \omega$, whence b must be a power of ω . If $p \neq 2$, noting that $\frac{1}{t_k} = t_k - 1$, we get

$$a = \frac{2(t_k + 1)}{2t_k\omega^j} = (t_k^2 - 1)\omega^{2j} = t_k\omega^{2j} = b\omega^j.$$

Direct computation shows that z^5 is scalar for the values given in the statement.

(iii) Assume that z^7 is scalar and note that $z^7e_3 = f_1e_1 + f_2e_2 + f_3e_3$, where

$$f_1 = a^3 - 3a^2b^2 + ab^4 + 4ab - b^3 - 1, \quad f_2 = 3a^2b - 4ab^3 - 2a + b^5 + 3b^2.$$

If $a = 0$, then $f_1 = -b^3 - 1$ and $f_2 = b^5 + 3b^2$. It follows that $p = 2$ and $b^3 = 1$. If $b = 0$, then $f_1 = a^3 - 1$ and $f_2 = 2a = 0$. It follows that $p = 2$ and $a^3 = 1$. So, assume $a, b \neq 0$. The resultant R of f_1, f_2 with respect to a (and to b) is:

$$(3) \quad R = t^{15} - 16t^{12} + 59t^9 - 67t^6 - 37t^3 + 8.$$

Now, we can factorize R as

$$(t^2 + t + 2)(t^3 - 2t^2 - t + 1)(t^4 - t^3 - t^2 - 2t + 4)(t^6 + 2t^5 + 5t^4 + 3t^2 + t + 1).$$

One can also verify that R splits over a field containing an element of order $\frac{21}{(p, 21)}$. $\square$

Lemma 3.4. *Assume that H is absolutely irreducible.*

- (i) *If $H \leq \text{SL}_3(q^2)$, then $H \leq \text{SU}_3(q^2)$ if, and only if, $b = a^q$.*
- (ii) *If q is odd and $H \leq \text{SL}_3(q)$, then $H \leq \text{SO}_3(q)$ if, and only if, $b = a$.*
- (iii) *Let $a = 0$. There exist no non-degenerate hermitian form and no non-degenerate orthogonal form fixed by H up to scalars.*

Proof. The characteristic polynomials of $z = xy$ and z^{-1} are respectively:

$$(4) \quad \chi_z(t) = t^3 - bt^2 + at - 1, \quad \chi_{z^{-1}}(t) = t^3 - at^2 + bt - 1.$$

The centralizers of x , y and z have respective dimensions 5, 3 and ≥ 3 . As $5+3+3 = 9+2$ and H is absolutely irreducible, by Scott's formula (see [11, Formula (2), page 324]), the centralizer of z must have the smallest possible dimension, i.e., 3. Thus, by [7, Theorem 3.16, page 207] z has a unique non-constant invariant factor (similarity invariant), namely its characteristic polynomial. Equivalently z is conjugate to a companion matrix. The same applies to z^{-1} and to z^σ .

(i) Let σ be as in (1). If $H \leq \text{SU}_3(q^2)$, then z^σ is conjugate to z^{-1} , whence $b = a^q$ from (4). Conversely, if $b = a^q$, from (4) and the considerations below it, we get that z^σ is conjugate to z^{-1} . By [11, Corollary 3.2, page 326] the group H fixes a non-degenerate Hermitian form, whence our claim.

(ii) If q is odd and $H \leq \text{SO}_3(q)$, then z is conjugate to z^{-1} , whence $b = a$. Conversely, if $b = a$, from (4) it follows that z is conjugate to z^{-1} . By the same Corollary quoted above, H fixes a non-degenerate bilinear form, whence our claim.

(iii) We argue by contradiction. Let J be a non-degenerate form fixed by H up to scalars, with $J^T = J^\tau$, $\tau \in \{\text{id}, \sigma\}$. From $x^T J x^\tau = \pm J$ and $\det(-I_3) = -1$ we get (I) $x^T J x^\tau = J$. It follows that J is non-scalar as $b \neq 0$. Moreover (II) $y^T J y = \omega^{\pm 1} J$, with $\omega \neq 1$ by (i), (ii) and the assumption $a = 0$, $b \neq 0$. In particular, $\omega \neq 1$ gives $p \neq 3$. Condition (II) forces $J = \text{diag}(1, \omega^j, \omega^{2j}) \cdot (c_1 y^0 + c_3 y + c_2 y^2)$. Condition $J^T = J^\tau$ gives $c_3 = \omega^{2j} c_2^\tau = \omega^{2j\tau} c_2^\tau$. Now $c_2 \neq 0$, as J is non-scalar, hence $\omega = \omega^\tau$. Thus

$$J = \begin{pmatrix} c_1 & c_2 & \omega^{\pm 2} c_2^\tau \\ c_2^\tau & \omega^{\pm 1} c_1 & \omega^{\pm 1} c_2 \\ \omega^{\pm 2} c_2 & \omega^{\pm 1} c_2^\tau & \omega^{\pm 2} c_1 \end{pmatrix}, \quad c_1 = c_1^\tau \in \mathbb{F}, \quad c_2 \in \mathbb{F}^*.$$

Now we impose that $D = x^T J x^\tau - J$ is zero. Assume first $c_1 = 0$. From $D_{2,3} = -2\omega^{\pm 1} c_2$ it follows $p = 2$. But, in this case, $D_{1,3} = c_2 b^\tau$ gives the contradiction $b = 0$. So we may suppose $c_1 = 1$. From $D_{2,3} = 0$ we get $b = -2c_2^\tau$, whence $p \neq 2$ and $c_2 = \frac{-b^\tau}{2}$. After this substitution, $D_{1,3} = 0$ gives:

$$-2\omega^{\pm 2} = \frac{b^{2\tau}}{b} = \frac{b^2}{b^\tau}.$$

Thus $b^{6\tau} = b^6 = -8b^3$ gives $b^3 = -8$. By Lemma 3.1(ii) H is reducible, a contradiction. $\square$

Theorem 3.5. *Let x, y be defined as in (2) with $a = 0$, $b \in \mathbb{F}_q^*$ such that:*

- (i) $b \neq -2\omega^j$, $\forall j \in \{0, 1, 2\}$;
- (ii) $\mathbb{F}_q = \mathbb{F}_p[b^3]$;
- (iii) $b \neq \omega^j$, $j \in \{0, 1, 2\}$ when $p = 2$.

Then $H = \text{SL}_3(q)$. Moreover for $q \geq 5$, there exists $b \in \mathbb{F}_q^*$ satisfying (i) to (iii) above. For $q = 2, 3$, taking $(a, b) = (1, 2)$ in (2), one gets $H = \text{SL}_3(q)$.

In particular, the groups $\text{SL}_3(q)$ and $\text{PSL}_3(q)$ are $(2, 3)$ -generated for all $q \neq 4$.

Proof. Let M be a maximal subgroup of $\text{SL}_3(q)$ which contains H . By Lemma 3.1 and assumption (i) the group H is absolutely irreducible. In particular M cannot belong to $\mathcal{C}_1 \cup \mathcal{C}_3$. Since $ab = 0$, by Lemma 3.2, the group M cannot belong to $\mathcal{C}_2$. By point (i) of Lemma 3.3 and Lemma 2.3 the group M cannot belong to $\mathcal{C}_6$. We

note that $z = xy$ has trace b . So, by assumption (ii), H is not conjugate to any subgroup of $\mathrm{SL}_3(q_0)Z$, with $q_0 < q$ and Z the center of $\mathrm{SL}_3(q)$. Thus M cannot belong to $\mathcal{C}_5$. If M is in class $\mathcal{S}$, then its projective image should be either $\mathrm{Alt}(6)$ or $\mathrm{PSL}_2(7)$. But this possibility cannot arise as, by (iii) and Lemma 3.3, the projective order of z is ≥ 8 . By point (iii) of Lemma 3.4 the group M does not belong to the class $\mathcal{C}_8$ and so we conclude that $H = \mathrm{SL}_3(q)$.

Now we come to the existence of b . If $q = p$, the elements $b \in \mathbb{F}_q^*$ which do not satisfy either (i) or (iii) are at most 3. Hence, for $q = p \geq 5$, there exists at least one $b \in \mathbb{F}_p^*$ satisfying conditions (i) and (ii). If $q \neq p$, it suffices to find an element $b \in \mathbb{F}_q^*$ satisfying (ii). This is possible for $q \neq 4$: indeed we may take b of order $q - 1$ since, by Lemma 2.4, $\mathbb{F}_p[b^3] = \mathbb{F}_p[b]$.

Finally let $q = 2, 3$, $(a, b) = (1, 2)$. The classes $\mathcal{C}_5$ and $\mathcal{S}$ are empty. Also the class $\mathcal{C}_8$ is empty for $q = 2$. For $q = 3$, assume that M is a subgroup of $\mathrm{SO}_3(3) \cong \mathrm{Sym}(4)$. By Lemma 3.3 this cannot happen, since the order of xy is greater than 4. We conclude that $H = \mathrm{SL}_3(q)$. $\square$

Theorem 3.6. *For $p = 2, 3, 5$ the groups $\mathrm{PSU}_3(p^2)$ are not $(2, 3)$ -generated.*

Proof. By what observed at the beginning of this Section, it is enough to show that, for $p = 2, 3, 5$, it is not possible to find a, b in the definition of x so that $H = \mathrm{SU}_3(p^2)$. By contradiction, suppose that a, b exist.

Point (i) of Lemma 3.4 gives $b = a^p$. Clearly $a^p \neq a$.

Case $q^2 = 25$. Set $\mathbb{F}_{25} = \{h + k\omega \mid h, k \in \mathbb{F}_5\}$. For $a \in \{\omega, -1 - \omega\}$, $ab = 1$ and H is monomial by Lemma 3.2. For $a \in \{2\omega, -2 - 2\omega\}$, H is reducible by point (i) of Lemma 3.1. For $a \in \{3\omega, -3 - 3\omega\}$, we have z^5 scalar, thus H has $\mathrm{Alt}(5)$ as composition factor. For $a \in \{\pm(3 + \omega), \pm(3 + 2\omega), \pm(1 + 3\omega)\}$ we have z^7 scalar and $[x, y]^4 = I$, thus H has $\mathrm{PSL}_2(7)$ as composition factor. For the remaining values, namely for $a \in \{-\omega, 1 + \omega, 1 + 2\omega, 1 - \omega, 2 + \omega, 3 - \omega, -1 + \omega, -1 + 3\omega\}$, the group H is reducible by point (ii) of Lemma 3.1.

Case $q^2 = 9$. Set $\mathbb{F}_9 = \mathbb{F}_3(\alpha)$, $\alpha^2 = \alpha + 1$. If $a \in \{\alpha, 2\alpha + 1\}$, then H is reducible. If $a \in \{\alpha + 1, \alpha^3 + 1\}$, then $ab = a^4 = 1$ and H is monomial. If $a \in \{-\alpha, -\alpha^3\}$, then $z = xy$ has order 7 and $[x, y]$ has order 4. It follows that $H \simeq \mathrm{PSL}_2(7)$.

Case $q^2 = 4$. Then $a = \omega^{\pm 1}$ and H is reducible by point (ii) of Lemma 3.1. $\square$

Theorem 3.7. *Suppose $q^2 \neq 4, 9, 25$. Define x, y as in (2) with $a \in \mathbb{F}_{q^2}^*$, $b = a^q$, and the further conditions:*

- (i) $a^q + a\omega^j + 2\omega^{2j} \neq 0$, $j \in \{0, 1, 2\}$;
- (ii) $a^{q+1} - 1 \neq 0$;
- (iii) $\mathbb{F}_{q^2} = \mathbb{F}_p[a^3]$;
- (iv) if $p \neq 2$, then $a^6 - 4a^3 - 1 \neq 0$;
- (v) $a^{15} - 16a^{12} + 59a^9 - 67a^6 - 37a^3 + 8 \neq 0$.

Then $H = \mathrm{SU}_3(q^2)$. Moreover there exists $a \in \mathbb{F}_{q^2}$ satisfying (i) to (v) above. In particular, $\mathrm{SU}_3(q^2)$ and $\mathrm{PSU}_3(q^2)$ are $(2, 3)$ -generated for all $q^2 \neq 4, 9, 25$.

Proof. By point (i) of Lemma 3.4 and the assumption $b = a^q$ we have that $H \leq \mathrm{SU}_3(q^2)$. Let M be a maximal subgroup of $\mathrm{SU}_3(q^2)$ which contains H . By Lemma 3.1 and conditions (i) and (iii), H is absolutely irreducible. In particular M cannot belong to $\mathcal{C}_1 \cup \mathcal{C}_3$. By (ii) we have $ab \neq 1$ so that M cannot belong to $\mathcal{C}_2$ by Lemma 3.2. Since z^6 is not scalar, by Lemma 3.3, we may apply Lemma 2.3 with $r = 3$ to deduce that M does belong to $\mathcal{C}_6$. From $b = a^q$ it follows $\mathbb{F}_p[b^3] = \mathbb{F}_p[a^3] = \mathbb{F}_{q^2}$.

As z has trace b , it cannot be conjugate to any element of $\mathrm{SL}_3(q_0)Z$, with $q_0 < q^2$ and Z the center of $\mathrm{SU}_3(q^2)$. Thus $M \notin \mathcal{C}_5$. Finally, if M is in class $\mathcal{S}$, then its projective image should be either $\mathrm{Alt}(6)$ or $\mathrm{PSL}_2(7)$, since we are excluding $q = 5$. But this possibility cannot arise as, by (iv), (v) and Lemma 3.3, the projective order of z is ≥ 8 . We conclude that $H = \mathrm{SU}_3(q^2)$.

As to the existence of some a satisfying all the assumptions, we first note that the elements of $\mathbb{F}_{q^2}$ which do not satisfy either (i) or (iv) or (v) are at most $3q + 6 + 15 = 3q + 21$. On the other hand, any $a \in \mathbb{F}_{q^2}^*$ of order $q^2 - 1$, satisfies (ii) and also (iii) by Lemma 2.4.

Assume first $q \geq 16$. By Lemma 2.2 there are at least $3q + 22$ elements $a \in \mathbb{F}_{q^2}^*$ having order $q^2 - 1$, and our claim is true.

For $q < 16$ we may take a of order $q^2 - 1$ and minimal polynomial specified in Table B. Indeed this minimal polynomial is coprime with each of $t^q + t\omega^j + 2\omega^{2j}$, for all j , $t^6 - 4t^3 - 1$ and $t^{15} - 16t^{12} + 59t^9 - 67t^6 - 37t^3 + 8$.

Table B

q	Minimal Polynomial	q	Minimal Polynomial
4	$t^4 + t + 1$	9	$t^4 + t^3 + t^2 + 2t + 2$
7	$t^2 + 6t + 3$	11	$t^2 + 7t + 2$
8	$t^6 + t^4 + t^3 + t + 1$	13	$t^2 - t + 2$

□

4. DIMENSION 5

In [15, Lemma 2.1, page 124] M. Vsemirnov parametrizes, up to conjugation, all the $(2, 3)$ irreducible pairs $X, Y \in \mathrm{SL}_5(\mathbb{F})$, for $p \neq 3$. His purpose is to show that $\mathrm{PSU}_5(4)$ is not $(2, 3)$ -generated. In his parametrization, taking $a = -1$, $d = 0$, $b, c \in \mathbb{F}$, we obtain

$$(5) \quad x = \begin{pmatrix} 0 & 1 & 0 & 0 & c \\ 1 & 0 & 0 & 0 & -c \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad c \neq 0, \quad y = \begin{pmatrix} 1 & 0 & -1 & 0 & b \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{pmatrix}.$$

Lemma 4.1. *Let x, y be as in (5). Then $H = \langle x, y \rangle$ is absolutely irreducible if, and only if, the following conditions hold:*

- (i) $b^2 + 3bc - b + 3c^2 - 3c + 1 \neq 0$, i.e., $b \neq (\omega^{\pm 1} - 1)c - \omega^{\pm 1}$;
- (ii) $b^2 + 10b + 16c + 9 \neq 0$, i.e., $(b, c) \neq (4\gamma - 1, -\gamma^2 - 2\gamma)$, $\forall \gamma \in \mathbb{F}$.

Proof. By Lemma 2.2 of [15], if $b^2 + 3bc - b + 3c^2 - 3c + 1 = 0$ the group H is reducible over $\mathbb{F}$. So condition (i) is necessary. If $(b, c) = (4\gamma - 1, -\gamma^2 - 2\gamma)$ for some $\gamma \in \mathbb{F}$, there is a 2-dimensional H -invariant space, namely $\langle w, yw \rangle$, where $w = (\gamma, -\gamma, 1, -1, 0)^T$. So condition (ii) is also necessary.

Now let W be a proper H -invariant subspace of $V = \mathbb{F}^5$. Denote by V_1^x , the eigenspace of x relative to 1. Notice that

$$xe_1 = e_2, \quad yxe_1 = e_3, \quad xyxe_1 = e_4, \quad (yx)^2e_1 = e_5.$$

Thus $e_1 \notin W$, since $W \neq V$. A direct calculation shows that $(I + y + y^2)V = \langle e_1 \rangle$. It follows that every element $w \in W$ satisfies the condition

$$(6) \quad w + yw + y^2w = 0.$$

Case 1 W is not contained in V_1^x . So there exists a non-zero vector $u \in W$ such that $w = u - xu \neq 0$. Calculation of $u - xu$ for a generic $u \in \mathbb{F}^5$ gives $w = (x_1, -x_1, x_3, -x_3, 0)^T$, for some $x_1, x_3 \in \mathbb{F}$. The vector w satisfies condition (6) if, and only if, $4x_1 - (b+1)x_3 = 0$. If $p \neq 2$, then we set $x_1 = \frac{b+1}{4}x_3$. Now, also xyw belongs to W and so it must satisfy condition (6). By condition (ii), we get the contradiction $w = 0$. If $p = 2$, we get $(b+1)x_3 = 0$. Condition (ii) implies $x_3 = 0$. In this case condition (6), applied to xyw , gives again $w = 0$.

Case 2 W is contained in V_1^x . Let $0 \neq w = (cx_1, cx_2, x_3, x_3, x_1 - x_2) \in W$. From $xyw = yw$ we get $x_3 = x_1 + (c-1)x_2$ whence $x_1(b+c) - x_2(b+c^2) = 0$, eliminating x_3 . If $b+c \neq 0$, we set $x_1 = \frac{b+c^2}{b+c}x_2$. Using condition (i) and the assumption $c \neq 0$, we see that w satisfies (6) only if it is zero, a contradiction. Finally assume $b = -c$. Then $xyw = yw$ gives $x_1 = x_3$ and $(c-1)x_2 = 0$. Moreover $xy^2w = y^2w$ gives $(c-1)x_1 = 0$. From $w \neq 0$ we conclude $c = 1$, $b = -1$, excluded by (i). $\square$

Lemma 4.2. *If H is absolutely irreducible, then it is not monomial.*

Proof. Assume that H acts monomially on a basis $B = \{v, \dots\}$. By the irreducibility of H , the permutations induced by x, y must generate a transitive subgroup of $\text{Sym}(5)$. In particular x must act as the product of two 2-cycles, y as a 3-cycle. This gives $p \neq 3$ as, for $p = 3$, the eigenspace of y relative to 1 has dimension 2. We may suppose that $xv = \pm v$. From $x^2 = 1$ and $\det(x) = 1$, we get $xv = v$, hence $v = (cx_1, cx_2, x_3, x_3, x_1 - x_2)^T$. It follows that:

$$B = \{v, yv, y^2v, xyv, xy^2v\}.$$

So B must be independent and, for some $j \in \{1, 2\}$, both vectors

$$(7) \quad y(xyv) - \omega^j(xyv), \quad y(xy^2v) - \omega^{-j}(xy^2v), \quad \omega \neq 1$$

have to be zero, since y must have the same similarity invariants of (5). Coordinates 4 and 5 of the vectors in (7) are 0 only if

$$x_3 = -\omega^{-j}cx_2, \quad x_1 = (2\omega^j c + 2c + 1)x_2.$$

After these substitutions, putting $x_2 = 1$, coordinates 2 and 3 of the two vectors are 0 only if $(b, c) = \left(-1, \frac{-\omega^j + 1}{3}\right)$. For $p = 2$, H is reducible by (ii) of Lemma 4.1. For p odd, coordinate 1 of the second vector in (7) is -2 , a contradiction. $\square$

Lemma 4.3. *Suppose H absolutely irreducible.*

- (i) *if q is odd and $H \leq \text{SL}_5(q)$, then $H \leq \text{SO}_5(q)$ if, and only if, $b = -1$;*
- (ii) *if $H \leq \text{SL}_5(q^2)$, then $H \leq \text{SU}_5(q^2)$ if, and only if, $b = c^q - c - 1$.*

Proof. The characteristic polynomials of $z = xy$ and z^{-1} are respectively

$$(8) \quad t^5 + t^4 + ct^3 + (-b - c - 1)t^2 - t - 1, \quad t^5 + t^4 + (b + c + 1)t^3 - ct^2 - t - 1.$$

The centralizers of x, y and z have respective dimensions 13, 9 and ≥ 5 . From $13 + 9 + 5 = 25 + 2$ and H absolutely irreducible, by Scott's formula we get that the centralizer of z must have dimension 5 (e.g. see [11]). Thus z has a unique non-constant invariant factor, namely its characteristic polynomial. Equivalently z is conjugate to a companion matrix. The same applies to z^{-1} and to z^7 , where

$\tau \in \{\text{id}, \sigma\}$. Thus, if $b = c^\tau - c - 1$, comparing the characteristic polynomials we get that z^τ is conjugate to z^{-1} . By [11, Corollary 3.2, page 326] the group H fixes a non-degenerate orthogonal or Hermitian form, whence our claim. $\square$

Lemma 4.4. *Assume H absolutely irreducible. Then*

- (i) *if $(xy)^k$ is scalar for some positive integer k , then $k \geq 10$;*
- (ii) *if $[x, y]^k$ is scalar for some positive integer k , then $k \geq 5$.*

Proof. (i) Let $D = (xy)^k$. For $k = 1, \dots, 5$, it suffices to look at the third row of D to conclude that D is not scalar. For $k = 6$, the condition $D_{3,1} = D_{3,2} = 0$ implies $b = -3$ and $c = 1$. So, it is easy to see that D is scalar only if $p = 2$, but in this case H is reducible. For $k = 7$, the condition $D_{3,1} = D_{3,2} = 0$ implies $(b, c) = (3, -1)$. However, in this cases D is not scalar. For $k = 8$, the condition $D_{3,1} - D_{5,1} = 0$ implies $(2c - 1)(b + 2c + 1) = 0$. If $p = 2$, then $b = 1$ and $D_{4,1} = 0$ implies $c = 0$. Assume $p \neq 2$. If $c = \frac{1}{2}$, then $D_{3,1} = 0$ implies $b = -\frac{9}{8}$. It is easy to see now that D is not scalar for these values of b, c . If $b = -2c - 1$, then the condition $D_{3,4} - D_{3,5} = 0$ implies $c = 2$. Also in this case one can verify that D is not scalar. For $k = 9$, the condition $D_{3,2} - D_{5,2} = 0$ implies $(b + 2c + 1)(2b - 3c^2 + 6c - 1) = 0$. If $b = -2c - 1$, then $D_{4,5} = 0$ implies $c^2 = -2$. In this case, it is easy to see that D is not scalar. Now, assume $2b - 3c^2 + 6c - 1 = 0$. If $p = 2$, this means $c = 1$. In this case $D_{3,1} = 0$ implies $b = 1$, but D is not scalar. If $p \neq 2$, take $b = \frac{3c^2 - 6c + 1}{2}$. We have $D_{3,1} = 3c^3 - \frac{11}{2}c^2 + 5c - \frac{5}{2} = 0$. If $p = 3$, we obtain $c^2 + c + 1 = 0$, i.e. $c = 1$. In this case H is reducible. So, assume $p \neq 3$. In this case $D_{3,1} = (3c - 3)(c^2 - \frac{5}{6}c + \frac{5}{6})$. It is now easy to see that D is not scalar.

(ii) Let $D = [x, y]^k$. For $k = 1, 2$, it suffices to look at the fifth row of D to conclude that D is not scalar. For $k = 3$, the condition $D_{3,1} - D_{5,1} = 0$ implies $c = 0$. For $k = 4$, the condition $D_{3,1} - D_{5,1} = 0$ implies $c^2 + 2b + 3c = 0$. If $p = 2$, we obtain $c = 1$. In this case, D is scalar only if $b = 1$. However, if $b = c = 1$, then H is reducible. Assume $p \neq 2$. Take $b = \frac{-c^2 - 3c}{2}$. The condition $D_{3,2} - D_{5,2} = 0$ implies $c = 1, -1$. In these cases D is not scalar. $\square$

Lemma 4.5. *Assume H absolutely irreducible. Then H is not contained in any subgroup $M = \langle \eta I \rangle \times S$, where S is isomorphic to $\text{PSU}_4(2)$ or $\text{PSL}_2(11)$.*

Proof. First, we observe that since H is absolutely irreducible, both x and y actually belong to the subgroup S , so $H \leq S$. Furthermore, by Lemma 4.4, we may assume that xy and $[x, y]$ have order at least 10 and 5, respectively. Also, we recall that $\text{Tr}(x) = 1$ and $\text{Tr}(y) = \text{Tr}(xy) = -1$.

Assume that S is isomorphic to $\text{PSU}_4(2)$. Then, $z = xy$ must be an element of order 12. This implies that $x \in 2B$ and $y \in \{3A, 3B\}$ in Atlas notation (simply, use the character table). On the other hand one can verify that in this case the order of $[x, y]$ is less than 5. This contradicts Lemma 4.4.

Assume that S is isomorphic to $\text{PSL}_2(11)$. Then, xy must be an element of order 11. We consider the character of an irreducible representation of degree 5 of $\text{PSL}_2(11)$ over $\mathbb{F}$. If $p \neq 2, 3, 11$, then $\chi(xy) = \frac{-1 \pm \sqrt{-11}}{2}$ which leads to $-2 = -1 \pm \sqrt{-11}$, i.e. $12 = 0$, in contradiction with the hypothesis on p . For $p = 2$, we have $\chi(xy) = \omega^{\pm 1}$, a contradiction. For $p = 3$ we have $\text{Tr}([x, y]) = 0 = b + 2c$. In this case, $z_{4,1} = c(c + 1)(c - 1)$. However, for $c = \pm 1$ the matrix $(xy)^{11}$ is not scalar. Finally, for $p = 11$ we obtain $\text{Tr}(xy) = -1 = \chi(xy) = 5$, a contradiction. $\square$

Theorem 4.6. *Let x, y as in (5) with $b = 0$ and $c \in \mathbb{F}_q^*$ such that:*

- (i) $3c^2 - 3c + 1 \neq 0$ and $16c + 9 \neq 0$;
- (ii) $\mathbb{F}_p[c^5] = \mathbb{F}_q$;
- (iii) if $p = 2$ and q is a square, then $c\sqrt{q} + c + 1 \neq 0$.

Then $H = \text{SL}_5(q)$. Moreover, if $q \neq 4, 16$, there exists $c \in \mathbb{F}_q^*$ satisfying (i), (ii), (iii). If $q = 4$, taking $b = c = \omega$ in (5), one gets $H = \text{SL}_5(4)$ and if $q = 16$, there exists $c \in \mathbb{F}_q^*$ such that $H = \text{SL}_5(16)$. In particular, the groups $\text{SL}_5(q)$ and $\text{PSL}_5(q)$ are $(2, 3)$ -generated for all q .

Proof. Let M be a maximal subgroup of $\text{SL}_5(q)$ which contains H . By Lemma 4.1 and condition (i), H is absolutely irreducible. In particular $M \notin \mathcal{C}_1 \cup \mathcal{C}_3$. Moreover $M \notin \mathcal{C}_2$ by Lemma 4.2. Since z^6 is not scalar by Lemma 4.4, we may apply Lemma 2.3 with $s = 5$ to deduce that $M \notin \mathcal{C}_6$. Since c is a coefficient of the characteristic polynomial of z (see (8)) and $\mathbb{F}_p[c^5] = \mathbb{F}_q$ by (ii), the matrix z cannot be conjugate to any matrix $(\lambda I)z_0$, with $(\lambda I) \in \text{SL}_5(\mathbb{F})$ and $z_0 \in \text{SL}_5(q_0)$, for any $q_0 < q$. Indeed, in this case, a coefficient of the characteristic polynomial of z_0 is $c\lambda^{-1} \in \mathbb{F}_{q_0}$, whence $c^5 = (c\lambda^{-1})^5 \in \mathbb{F}_{q_0}$, in contrast with (ii). We conclude that M does not belong to class $\mathcal{C}_5$.

We now exclude that $M \in \mathcal{C}_8$. Let J be a non singular matrix such that $x^T J x^\tau = \lambda J$ and $y^T J y^\tau = \mu J$, for some $\lambda, \mu \neq 0$. Considering the orders of x, y we get $\lambda^2 = \mu^3 = 1$. On the other hand, considering the determinants, we obtain $\lambda^5 = \mu^5 = 1$. We conclude that $\lambda = \mu = 1$. In other words, if H fixes J up to scalars, then it actually fixes J . By Lemma 4.3(i) H does not fix any orthogonal form. On the other hand, assume that q is a square and that $H \leq \text{SU}_5(q)$. For $b = 0$, Lemma 4.3(ii) gives $c\sqrt{q} = c + 1$. It follows $c = c^q = (c + 1)\sqrt{q} = c + 2$, whence $p = 2$. So this case is excluded by assumption (iii).

Finally, suppose M in class $\mathcal{S}$. By Lemma 4.5 we have to consider only the case $q = 3$ and $M = M_{11}$. However, if $c = \pm 1$, then xy has order, respectively, 16 and 121, a contradiction. We conclude that $H = \text{SL}_5(q)$.

As to the existence of some c satisfying all the assumptions in the statement, the elements of $\mathbb{F}_q^*$ which do not satisfy (i) are at most 3. Moreover, by Lemma 2.4, if $q \neq 16$, any element in $\mathbb{F}_q^*$ of order $q - 1$ satisfies (ii). Hence, if p is odd and $q \geq 9$, there exists c of order $q - 1$ satisfying all conditions since $\varphi(q - 1) > 3$. If $q = 3, 5, 7$, one can check that $c = -2$ satisfies (i) and (ii).

Now assume $p = 2$. If q is not a square, any element in $\mathbb{F}_q^*$ of order $q - 1$ satisfies all the assumptions. If q is a square, assume first $q \neq 2^2, 4^2$. The elements in $\mathbb{F}_q^*$ of order $q - 1$, which do not satisfy either (i) or (iii), are at most $\sqrt{q} + 2$. Moreover all elements of order $q - 1$ satisfy (ii). We have $\varphi(q - 1) > 4\sqrt{q} > \sqrt{q} + 2$ by Corollary 2.2 and direct calculation if $q < 16^2$. Thus c exists if $q \neq 2^2, 4^2$ and so $H = \text{SL}_5(q)$. When $q = 4^2$, we take c of order 15, with minimal polynomial $t^4 + t^3 + 1$. Then c satisfies (i) and (iii). Moreover $z = xy$ has order divisible by 41, a prime which does not divide $|\text{SL}_5(2^m)|$ for any $m \leq 3$. So $H = \text{SL}_5(4^2)$.

When $q = 2^2$, we take $b = c = \omega$. In this case, the classes $\mathcal{C}_6$ and $\mathcal{S}$ are empty. By Lemmas 4.1 and 4.2 the subgroup H is irreducible and not monomial. Furthermore, $\mathbb{F}_4 = \mathbb{F}_2(\omega)$ and by Lemma 4.3 H does not fix non-degenerate forms. Again $H = \text{SL}_5(2^2)$. $\square$

We recall that the group $\text{PSU}_5(4)$ is not $(2, 3)$ -generated, see [15].

Theorem 4.7. *Let x, y be as in (5) with $c \in \mathbb{F}_{q^2}^*$, $b = c^q - c - 1$ and suppose:*

- (i) $c^{2q} + c^{q+1} - 3c^q + c^2 - 3c + 3 \neq 0$;
- (ii) $c^{2q-1} - 2c^q + 8c^{q-1} + c + 8 \neq 0$;
- (iii) $\mathbb{F}_{q^2} = \mathbb{F}_p[c^5]$.

Then $H = \mathrm{SU}_5(q^2)$. Moreover, if $q^2 \neq 2^2, 4^2$, there exists $c \in \mathbb{F}_{q^2}^*$ satisfying conditions (i) to (iii). If $q^2 = 4^2$, there exists $c \in \mathbb{F}_{q^2}^*$ such that $H = \mathrm{SU}_5(16)$. In particular, the groups $\mathrm{SU}_5(q^2)$ and $\mathrm{PSU}_5(q^2)$ are $(2, 3)$ -generated for all $q^2 \geq 9$.

Proof. $H \leq \mathrm{SU}_5(q^2)$ by Lemma 4.3 and the assumption $b = c^q - c - 1$. Let M be a maximal subgroup of $\mathrm{SU}_5(q^2)$ which contains H . Conditions (i) and (ii) correspond to those in Lemma 4.1, setting $b = c^q - c - 1$. Thus H is absolutely irreducible. In particular $M \notin \mathcal{C}_1 \cup \mathcal{C}_3$. Moreover M cannot belong to $\mathcal{C}_2$ by (iii) and Lemma 4.2. Since z^6 is not scalar by Lemma 4.4, we may apply Lemma 2.3 with $s = 5$ to deduce that $M \notin \mathcal{C}_6$. We have $\mathbb{F}_p[c^{5q}] = \mathbb{F}_p[c^5] = \mathbb{F}_{q^2}$, by (iii). Since $b + c + 1 = c^q$ is a coefficient of the characteristic polynomial of z (see (8)), the matrix z cannot be conjugate to any element of $\mathrm{SL}_5(q_0)Z$, with $q_0 < q^2$ and Z the center of $\mathrm{SU}_5(q^2)$. Thus $M \notin \mathcal{C}_5$. Finally, M cannot be in class $\mathcal{S}$ by Lemma 4.5. We conclude that $H = \mathrm{SU}_5(q^2)$.

As to the existence of some c satisfying all the assumptions, suppose $q^2 \neq 16$. Any element of $\mathbb{F}_{q^2}^*$ of order $q^2 - 1$ satisfies (iii) by Lemma 2.4. The non-zero elements in $\mathbb{F}_{q^2}$ which do not satisfy either (i) or (ii) are at most $2q + 2q - 1 = 4q - 1$. If $q \geq 16$, by Lemma 2.2 there are at least $4q$ elements in $\mathbb{F}_{q^2}^*$ having order $q^2 - 1$. So c exists. For $q = 4, 7, 8, 9, 11, 13$, we may take c with minimal polynomial m_c as in Table B and for $q = 3, 5$ we take c of order $q^2 - 1$, with minimal polynomial $m_c = t^2 - t - 1$ and $m_c = t^2 - t + 2$, respectively. Then c satisfies (i) and (ii) since m_c is coprime with

$$p_1 = t^{2q} + t^{q+1} - 3t^q + t^2 - 3t + 3, \quad p_2 = t^{2q-1} - 2t^q + 8t^{q-1} + t + 8.$$

It also satisfies (iii) except for $q = 4$. In this case, z has order divisible by 17, a prime which does not divide $|\mathrm{SU}_5(2^2)|$. Thus $H = \mathrm{SU}_5(q^2)$, for all $q \neq 2$. $\square$

REFERENCES

- [1] Bray, J., Holt, D., Roney-Dougal, C., The Maximal Subgroups of the Low-Dimensional Finite Classical Groups, London Math. Soc. Lecture Note Series 407, Cambridge University Press (2013).
- [2] Carter R.W., Simple groups of Lie type, Pure and Applied Mathematics, Vol. 28. John Wiley & Sons, London-New York-Sydney, 1972
- [3] Cohen, J., On non-Hurwitz groups and non-congruence subgroups of the modular group, Glasgow Math. J. 22 (1981), 1–7.
- [4] Di Martino, L., Vavilov, N., $(2, 3)$ -generation of $\mathrm{SL}_n(q)$, I, Cases $n = 5, 6, 7$, Comm. Algebra, 22, (1994), 1321–1347.
- [5] Fricke, R., Klein, F., Vorlesungen über die Theorie der Elliptischen Modulfunctionen, 2 Vols., Teubner, Leipzig (1890-2).
- [6] Garbe, D., Über eine classe von arithmetisch definierbaren Normalteilern der Modulgruppe, Ann. Math., 235 (1978), No. 3, 195–215.
- [7] Jacobson N., Basic algebra. I. Second edition, W. H. Freeman and Company, New York, 1985.
- [8] Liebeck, M.W., Shalev, A., Classical groups, probabilistic methods, and the $(2, 3)$ -generation problem, Ann. Math. (2) 144 (1996), 77–125.
- [9] Macbeath, A.M., Generators for the linear fractional groups, Proc. Simp. Pure Math. 12 (1969), 14–32.

- [10] Mazurov, V., Khukhro, E., Kourovka notebook No.18, Unsolved problems in Group Theory, arXiv 1401.0300.
- [11] Pellegrini, M.A., Tamburini Bellani, M.C., Vsemirnov, M.A., Uniform $(2, k)$ -generation of the 4-dimensional classical groups, J. Algebra, 369 (2012), 322–350.
- [12] Tabakov, K., $(2, 3)$ -generation of the groups $\mathrm{PSL}_7(q)$, Proceedings of the Forty Second Spring Conference of the Union of Bulgarian Mathematicians, Borovetz, April 2–6, 2013.
- [13] Tabakov, K., Tchakerian, K., $(2, 3)$ -generation of the groups $\mathrm{PSL}_6(q)$, Serdica Math J. 37 (2011) No. 4, 365–370.
- [14] Tchakerian, K., $(2, 3)$ -generation of the groups $\mathrm{PSL}_5(q)$, Ann. Univ. Sofia, Fac. Math. Inf., 97 (2005), 105–108.
- [15] Vsemirnov, M., More classical groups which are not $(2, 3)$ -generated, Arch. Math. (Basel) 96 (2011), no. 2, 123–129.
- [16] Vsemirnov, M., On $(2, 3)$ -generated groups, Group Theory Conference in honour of V. Mazurov, Novosibirsk, July 2013.
- [17] Wagner, A., The minimal number of involutions generating some finite three dimensional groups, Boll. Un. Mat. Ital. (5) 15-A (1978), 431–439.

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ CATTOLICA DEL SACRO CUORE, VIA MUSEI 41, I-25121 BRESCIA, ITALY

E-mail address: marcoantonio.pellegrini@unicatt.it

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ CATTOLICA DEL SACRO CUORE, VIA MUSEI 41, I-25121 BRESCIA, ITALY

E-mail address: mariaclara.tamburini@unicatt.it